

Shadow features and shadow bands in the paramagnetic state of cuprate superconductors

Y. M. Vilk

Materials Science Division, Argonne National Laboratory,
Argonne, IL 60439
e-mail: yvilk@hexi.msd.anl.gov

The conditions for the precursors of antiferromagnetic bands in cuprate superconductors are studied using weak-to-intermediate coupling approach. It is shown that there are, in fact, three different precursor effects due to the proximity to antiferromagnetic instability: i) the shadow band which associated with new pole in the Green's function ii) the dispersive shadow feature due to the thermal enhancement of the scattering rate and iii) the non-dispersive shadow feature due to quantum spin fluctuation that exist only in $\vec{k}$ -scan of the spectral function $A(\omega_{Fixed}, \vec{k})$. I found that dispersive shadow peaks in $A(\omega, \vec{k})$ can exist at *finite* temperature T in the renormalized classical regime, when $T \gg \omega_{sf}$, $\xi_{AFM} > \xi_{th} = v_F/T$ (ω_{sf} is the characteristic energy of spin fluctuations, ξ_{th} is the thermal wave length of electron). In contrast at zero temperature, only non-dispersive shadow feature in $A(\omega_{Fixed}, \vec{k})$ has been found. I found, however, that the latter effect is always very small. The theory predict no shadow effects in the optimally doped materials. The conditions for which shadow peaks can be observed in photoemission are discussed.

PACS numbers: 71.27,79.60

I. INTRODUCTION

Recent advances in the Angular Resolved Photoemission technique (ARPES) have allowed to determine electronic excitations of some High- T_c materials. Two different approaches to photoemission are currently used. In type I ARPES one measures $A(\omega, \vec{k}_{Fixed})$ the probability that an excitation of fixed momentum $\vec{k}_{Fixed}$ has an energy ω . In the type II ARPES one measures the probability $A(\omega_{Fixed}, \vec{k})$ that an excitation of fixed energy ω_{Fixed} has a momentum $\vec{k}$. Using the latter technique on the high-temperature superconductor B2212, Aebi et al.¹ have found, in addition to the main Fermi surface (F. S.), another F. S. that is identical to the first one but of much smaller intensity and displaced from the main F. S. by the antiferromagnetic (AFM) wave vector $\vec{Q} = (\pi/a, \pi/a)$. This structure has been taken as evidence for the existence of the "shadow bands" that had been predicted much earlier by Kampf and Schrieffer³. These authors had argued that precursors of the AFM bands should appear in the paramagnetic state close to the AFM instability. More recently, observation of dispersing shadow bands has been also reported in type I photoemission experiments². However, the AFM interpretation of the phenomenon observed in B2212 remains controversial because: i) the AFM correlation length ξ in optimally doped B2212 is rather short⁴, ii) the effect has not been found yet in any other high-temperature superconductor or in the underdoped B2212 iii) it becomes more pronounced when one increase orthorombicity by lead doping⁵. All these leave open the possibility of structural origin of the phenomenon observed in Refs.^{1,2}. It is thus very important to understand under which general conditions shadow bands can occur in prac-

tice. Some aspects of this issue have been considered in a number of recent theoretical publications⁶⁻⁹. In particular, the strong coupling numerical calculations for the t-J model⁹ show peaks outside non-interacting F. S. in the underdoped case for $T = 0$. However, these results have been obtained so far only for very small clusters and the results close to optimal doping are unclear⁹.

In this paper, I study under which general conditions shadow bands can occur, using a well known method that can be justified up to intermediate coupling. I show that there are, in fact, a few different precursors effects due to the proximity to AFM instability depending on whether the effect is due to classical $\omega_{sf} \ll T$ or quantum $T \ll \omega_{sf}$ spin fluctuations (here ω_{sf} is characteristic energy of spin fluctuations). I also show that shadow effects in type I and type II photoemission experiments may occur under different conditions. For type I ARPES experiments, I find that while classical spin fluctuations can produce dispersive shadow effects in $A(\omega, \vec{k})$, quantum spin fluctuations cannot. The key difference between both regimes is that classical thermal fluctuations lead to precursors of the magnetic Bragg peaks in the *static* spin structure factor, while the dynamical nature of quantum spin fluctuations does not allow sharp structures in $S_{sp}(\vec{q})$, no matter how large ξ/a is (a lattice constant). Consequently, shadow effects will appear in type I experiments at low temperature only if the AFM correlation length ξ has the appropriate temperature dependence, not just when ξ/a is large enough. This conclusion is qualitatively different from the results of numerical studies⁷⁻⁹, in which it was tacitly assumed that the effect is determined by the parameter ξ/a alone. It is only in type II ARPES experiments that quantum spin fluctuations can lead to a non-dispersive shadow feature

in $A(\omega_{Fixed}, \vec{k})^{10}$. I find, however, that the latter effect is very small and exists only in a narrow range near zero energy. More experimental implications of my results are discussed in the conclusion.

II. MODEL AND CALCULATIONAL PROCEDURE

I evaluate the effect of spin fluctuations on the electronic self-energy in the one-loop approximation. The expression for the self-energy has the form:

$$\Sigma(\omega, \vec{k}) = g\bar{g} \int \frac{d^2 q d\omega'}{(2\pi)^3} \chi_{sp}'(\omega', \vec{q}) \left[\coth\left(\frac{\omega'}{2T}\right) - \tanh\left(\frac{\varepsilon(\vec{k}+\vec{q})}{2T}\right) \right] \times \frac{1}{\omega - \omega' - \varepsilon(\vec{k}+\vec{q}) + i0} \quad (1)$$

where $\chi_{sp}(\omega, \vec{q})$ is the spin susceptibility, $g\bar{g}$ is the effective coupling constant between electrons and spin fluctuations and $\varepsilon(\vec{k})$ is the energy dispersion relative to the chemical potential. The constant vertex corrections are included in $\bar{g}$. It was shown within the Hubbard model⁶, that by including vertex corrections in this way the above self-energy expression (1) remains a good approximation up to intermediate coupling.

Close to AFM instability the low-energy asymptotic form of the spin susceptibility χ_{sp} can be written in the Orstein-Zhernike form:

$$\chi_{sp}(\omega, \vec{q}) = \frac{\chi_{sp}(0, \vec{Q})}{1 + (\vec{q} - \vec{Q})^2 \xi^2 - i\omega/\omega_{sf}} \quad (2)$$

where ξ is the AFM correlation length and ω_{sf} is the characteristic energy of the spin fluctuations. Any RPA-like theory, whether it uses bare or renormalized vertices⁶ or propagators⁸, have the Orstein-Zhernike form close to $\vec{q} = \vec{Q}$. In the context of High- T_c materials, the Orstein-Zhernike susceptibility with particular set of parameters is often called MMP susceptibility. It has been used to fit a number of experiments, that are sensitive to the strong enhancement of χ around $\vec{q} = \vec{Q}^{11}$. Here, I will adopt a more general approach and consider for which general conditions on the parameters ξ and ω_{sf} the precursors of AFM bands can exist. Since shadow effects come from the peak in χ around $\vec{q} = \vec{Q}$, the Orstein-Zhernike form of χ is sufficient for my purposes and its simplicity will allow me to obtain a number of results analytically. The imaginary part of the susceptibility $\chi_{sp}''(\omega, \vec{q})$ has to be cut off at some high frequency Γ_{cut} in order to satisfy the local moment sum rule. In electronic models this cutoff is of the order of a few hopping integrals t . For the low energy physics the precise value of Γ_{cut} is not important. I use $\Gamma_{cut} = 5.5t$.

From a microscopic point of view (see, for example,¹¹), the parameters in the susceptibility Eq.(2) scale with the correlation length as: $\chi_{sp}(\vec{Q}, 0) \propto \xi^2$, $\omega_{sf} \propto 1/\xi^2$.

Consequently the model is defined by only three parameters ξ , ω_{sf} and $g' \equiv g\bar{g}\chi_{sp}(\vec{Q})/\xi^2$. Qualitatively, our results mainly depend on ξ and on T/ω_{sf} while the parameter g' (in the Hubbard model⁶ $g' \propto U$) determines the quantitative scale of the effects. To model the band-structure of the hole-doped high- T_c materials I use the simplest tight-binding model that correctly reproduces the experimentally observed Fermi surfaces in YBCO and B2212 at optimal doping, namely $\epsilon(\vec{k}) = -2t(\cos(k_x) + \cos(k_y)) - 4t'\cos(k_x)\cos(k_y)$ with $t' = -0.45t$ (I set $a = 1$). The band structure estimate for t is about $t \approx 250meV$. Using this approach, I first present numerical results and then analytically find the conditions for the existence of the various effects.

Both the self-energy and the spectral function $A(\omega, \vec{k}) \equiv -2\Sigma''(\omega, \vec{k})/[(\omega - \epsilon(\vec{k}) + \mu - \Sigma'(\omega, \vec{k}))^2 + |\Sigma''(\omega, \vec{k})|^2]$ are discussed. The chemical potential of the interacting electrons is found from the usual condition $\int f(\omega)A(\omega, \vec{k})d\omega d^2k/(2\pi)^3 = n/2$. I choose $n = 0.83$ that corresponds to optimal doping in B2212¹².

III. RESULTS

I start with numerical results and figures that illustrate various points I wish to make. Analytical results for various regimes identified in the numerical examples appear in a subsequent section.

A. Illustrative examples

The important aspect of critical AFM fluctuations is that they lead to strongly $\vec{k}$ -dependent self-energy. The maximum of the scattering rate $-2\Sigma''(0, \vec{k})$ at finite temperature occurs on the so-called shadow Fermi surface (Sh. F. S.). The latter is *defined* here as the set of points obtained by the translation of the real F. S. by the wave vector $\vec{Q} = (\pi, \pi)$. The inset in Fig.1(a) shows the real F.S. and the Sh. F. S. for the optimally doped materials. The points at which the real F. S. and the Sh. F. S. intersect are the points at which the spectral function $A(\omega, \vec{k}_F)$ is affected the most by spin fluctuations. I start the analysis with these crossing points, $\vec{k} = \vec{k}_{cr}$, also known as hot spots.

1. Shadow bands at the hot spots

The filled line on Fig.1 shows the spectral function $A(\omega, \vec{k}_{cr})$ in the regime dominated by classical thermal fluctuations ($\omega_{sf} \ll T$, $T = 0.03t$ and $\xi = 20a$). We see that instead of the usual quasi-particle peak at $\omega = 0$ there are two precursors of the antiferromagnetic bands with a pseudogap between them. These bands are associated with two new quasiparticle solutions of the equation

$\omega - \epsilon(\vec{k}) + \mu - \Sigma'(\omega, \vec{k}) = 0$. It is because these solutions exist that the term *bands* is used. I also find that in this case the scattering rate $\Sigma''(\omega, \vec{k}_{cr})$ has maximum at the Fermi level $\omega = 0$.

I will show now by contradiction, that this effect exists only if $\xi(T)$ increases sufficiently rapidly with decreasing temperature. Indeed, suppose the correlation length does not increase when temperature decreases below the temperature $T = 0.03$ used for the filled line on Fig.1. Taking a much smaller temperature $T/t = 0.005$ while keeping all other parameters fixed, the dashed line on Fig.1 shows that the pseudogap in $A(\omega, \vec{k}_{cr})$ disappears while the quasiparticle peak reappears. I find similar results when I keep the temperature and ξ constant but increase ω_{sf} up to $\omega_{sf} \approx T$. This means that the pseudogap in $A(\omega, \vec{k}_{cr})$ can exist only if the correlation length $\xi(T)$ increases sufficiently rapidly at low temperature. I show below that $\xi(T) \propto \exp(\text{const}/T)$ is a sufficient condition. The latter condition is realized in the renormalized classical regime (R. C.) $\omega_{sf} \propto \xi^{-2} \ll T$ that always precedes the $T = 0$ phase transition in two dimensions.

2. Shadow features in the diagonal direction $(0,0) - (\pi,\pi)$

In this direction, real and shadow F.S. are well separated and $|\epsilon(\vec{k}) - \epsilon(\vec{k} + \vec{Q})|$ is large. The filled curve in Fig.2 shows the spectral function at $T = 0.03t$ for a wave vector $\vec{k} = (0.66, 0.66)\pi$ that is just a little bit inside the Sh. F. S. ($\vec{k}_{Sh.F} \simeq (0.63, 0.63)\pi$). The quasiparticle peak in this case is at large positive energies and cannot be seen in the photoemission experiments since they measure $f(\omega)A(\omega, \vec{k})$. There is no new quasiparticle solution to $\omega - \epsilon(\vec{k}) + \mu - \Sigma'(\omega, \vec{k}) = 0$ in this case. Nevertheless, there is a local maximum at negative binding energies (shown by arrow) that coincides with the maximum in the imaginary part, $\Sigma''(\omega, \vec{k})$ in Fig. 2b. I call this a *shadow feature*. This maximum occurs because for large values of $|\omega - \epsilon(\vec{k}) + \mu|$, the quantity $\omega - \epsilon(\vec{k}) + \mu - \Sigma'(\omega, \vec{k})$ is large and slowly varying in the region where $\Sigma''(\omega, \vec{k})$ acquires a maximum. Hence, this maximum is directly reflected in the spectral function that becomes proportional to $\Sigma''(\omega, \vec{k})$ rather than inversely proportional to it as in the case of quasiparticles. This maximum occurs at $\omega \approx \tilde{\epsilon}(\vec{k} + \vec{Q})$, a dispersion that looks like the AFM band but with the gap equal to zero. Nevertheless, this is definitely *not* a real AFM band since a) it is thermally excited (see below) b) it exists only in certain regions of $\vec{k}$ -space where $|\epsilon(\vec{k}) - \epsilon(\vec{k} + \vec{Q})| \gg \Sigma''(\omega, \vec{k})$. Note that similar shadow features have been found in the Hubbard model with $t' = 0$, using the FLEX approximation⁸.

To show that this shadow feature is a thermally excited state I plot the dashed line in Fig.2 for the case where the temperature is decreased to $T = 0.003t$ with all other parameters the same as for $T = 0.03t$. Clearly

the shadow feature (shown by arrow) disappears at lower temperature. It also disappears for $\omega_{sf} > T$. I note that the maximum in $A(\omega, \vec{k})$ around $\omega_{max} \sim -t$ is different from the “shadow band” observed in Ref.². It stays at $\omega_{max} \sim -t < 0$ even when $\vec{k}$ crosses the Sh. F. S..

3. Non-dispersive $T \rightarrow 0$ shadow feature in type II configuration:

The above considerations lead to the conclusion that only classical thermal spin fluctuations that destroy long range order at $T \neq 0$ in two dimensions can produce the above dispersive shadow peaks in both type I and type II experiments. Quantum spin fluctuations $T \ll \omega_{sf}$ alone cannot produce shadow effect in $A(\omega, \vec{k}_{Fixed})$ measured in type I ARPES, no matter how large is ξ/a . However, in the type II configuration, ($\vec{k}$ -dependence of $A(\omega_{Fixed}, \vec{k})$), quantum spin fluctuations can lead to a non-dispersive shadow feature at $T = 0$ when $\xi/a \gg 1$ (Ref.¹⁰). The effect is due to the enhancement of $\Sigma''(\omega_{Fixed}, \vec{k})$ on the Sh.F.S. at finite ω . For small ω the maximum in $\Sigma''(\omega_{Fixed}, \vec{k})$ translates into a maximum in $A(\omega_{Fixed}, \vec{k})$ at $\vec{k} = k_{Sh.F.S.}$, since $A(\omega_{Fixed}, \vec{k}) \propto \Sigma''(\omega_{Fixed}, \vec{k})$ for $\vec{k}$ far from the F.S.. Since $A(0, \vec{k} \neq \vec{k}_{F.S.}) = 0$ at $T = 0$, one needs to increase ω to have a noticeable weight for this maximum in the incoherent background. However, when ω increases the width of the maximum in k -space rapidly increases and above a certain energy $\omega_0 < t$, mainly determined by $(\xi/a)^2 \omega_{sf}$, the maximum disappears. This occurs because the effect from Σ'' is offset by the contribution from the denominator in $A(\omega, \vec{k}) \approx \Sigma''(\omega, \vec{k})/(\omega - \tilde{\epsilon}(\vec{k}))^2$. For $|\omega| > \omega_0$, the closer $\vec{k}$ is to the real F. S. the larger $A(\omega, \vec{k})$ is. This is illustrated in the inset of Fig.2. I found that the effect is always very small even for large values of ξ and $\xi^2 \omega_{sf}$. For values of $\xi = 2.3a$ and $\omega_{sf} \approx 10meV$ which are typical for cuprates closed to the optimal doping¹¹ it does not exist even at very small $\omega \ll t$.

B. Analytical results

1. Difference between classical and quantum regimes

I now explain analytically why the effect of classical and quantum AFM fluctuation on $A(\omega, \vec{k})$ are qualitatively different. When $\omega_{sf} \ll T$ the effect of spin fluctuations on single-particle properties can be considered quasistatic. Indeed, neglecting ω' in the energy denominator of the self-energy formula Eq. (1) one obtains¹³:

$$\Sigma(\omega, \vec{k}) = g\bar{g} \int \frac{d^2q}{(2\pi)^2} \frac{S_{sp}(\vec{q})}{\omega - \tilde{\epsilon}(\vec{k} + \vec{q}) + i0} \quad (3)$$

where $S_{sp}(\vec{q})$ is the static structure factor $S_{sp}(\vec{q}) = \int \chi''_{sp}(\omega', \vec{q}) \coth(\omega'/2T) d\omega'/2\pi$. The key point is that in

the RC regime, $S_{sp}(\vec{q})$ has the *same* singular behavior as the static susceptibility $\chi'_{sp}(0, \vec{q})$ because for $q \approx Q$, $S_{sp}(\vec{q}) \approx T \int (d\omega/\pi) \chi''(\omega, \vec{q})/\omega = T \chi'_{sp}(0, \vec{q})$. Since close to the AFM instability $\chi'_{sp}(0, \vec{q})$ is a Lorentzian peaked at $q = Q$, the static structure factor $S_{sp}(\vec{q})$ has a Lorentzian maximum in the regime $\omega_{sf} \ll T$:

$$S_{sp}(\vec{q}) \propto \frac{T}{\xi^{-2} + (\vec{q} - \vec{Q})^2} \quad (4)$$

The presence of a sharp precursor of the Bragg peak in the RC regime leads to the shadow peak in $A(\omega, \vec{k}_{Fixed})$. The situation is qualitatively different in the case $T \ll \omega_{sf}$ and $\xi/a \gg 1$. In that case, $S_{sp}(\vec{q}) \approx \int_0^\infty \chi''(\omega, \vec{q}) d\omega/\pi$ and using Eq. (2) I obtain:

$$S_{sp}(\vec{q})|_{T=0} \propto \ln[\xi^{-2} + (\vec{q} - \vec{Q})^2] \quad (5)$$

The logarithmic singularity of $S_{sp}(\vec{q})$ in the case $T \ll \omega_{sf}$ is integrable and does not lead to any divergence in $\Sigma(\omega, \vec{k})$ when $\xi \rightarrow \infty$. Consequently, there is no shadow effects in $A(\omega, \vec{k}_{Fixed})$, when electrons approach the AFM instability $\xi \gg a$ at zero temperature $T = 0$. This conclusion is different from the result of Kampf and Schrieffer³ who used the zero temperature formalism and found shadow bands for $\xi \approx 20a$. The difference is due to the fact that the phenomenological susceptibility used in Ref.³ was separable in $\vec{q}$ and ω , $\chi_{sp, KSh} = f(\vec{q})g(\omega)$. In that case, the $\vec{q}$ -dependence of the static structure factor $S_{sp}(\vec{q})$ is *always* the same as that of the susceptibility $\chi'_{sp}(0, \vec{q})$. Consequently, the static structure factor $S_{sp, KSh}(\vec{q})$ has a Lorentzian-like peak even when $T \ll \omega_{sf}$. The separability in ω and $\vec{q}$ of the susceptibility is an oversimplification that can be partially justified only in the regime $\omega_{sf} \ll T$. Thus, the results of Ref.³ should not be applied to the zero-temperature paramagnetic state.

2. Conditions on $\xi(T)$

The asymptotic form of $\Sigma(\omega, \vec{k})$ in the RC regime has been obtained in Ref.⁶ for the case $n = 1$, $t' = 0$. The generalization to the present case is straightforward and for Σ'' I get, for $\xi > \xi_{th} = v_{\vec{k}+\vec{Q}}/T$:

$$\Sigma''(\omega, \vec{k}) = -\frac{g'T}{4\sqrt{[\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})]^2 + v_{\vec{k}+\vec{Q}}^2 \xi^{-2}}} \quad (6)$$

We see that $\Sigma''(\omega, \vec{k})$ has a maximum when $\omega = \tilde{\varepsilon}(\vec{k} + \vec{Q})$ with $\Sigma''(\tilde{\varepsilon}(\vec{k} + \vec{Q}), \vec{k}) = -g'T\xi/4v_{\vec{k}+\vec{Q}}$. The effect is largest at the Van Hove point $v_k = 0$, in which case $\Sigma''(\tilde{\varepsilon}(\vec{k} + \vec{Q}), \vec{k}) \propto \xi^2$. Since away from $k = k_{F.S.}$, we have $A(\omega, \vec{k}) \propto \Sigma''(\omega, \vec{k})$, we obtain for $\xi > \xi_{th}$, the shadow feature discussed above. On the other hand, at the crossing points $\tilde{\varepsilon}(\vec{k}_{cr} + \vec{Q}) = \tilde{\varepsilon}(\vec{k}_{cr}) = 0$, $A(0, \vec{k}) \propto 1/\Sigma''(0, \vec{k})$

and the enhancement in Σ'' leads to the AFM pseudogap with no quasiparticle peak inside. This is similar to what was found in the case of perfect nesting ($n = 1$, $t' = 0$)⁶, except that in the latter case the pseudogap opens up over all the F.S., since the real F.S. and the Sh.F.S coincide (all points on the F.S. are crossing points).

The real part of the self-energy can be obtained from Eq.(6) using Kramers-Kronig relation and has the form:

$$\Sigma'(\omega, \vec{k}) = \frac{g'T}{4\pi\sqrt{[\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})]^2 + v_{\vec{k}+\vec{Q}}^2 \xi^{-2}}} \times \ln \left| \frac{\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q}) + \sqrt{[\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})]^2 + v_{\vec{k}+\vec{Q}}^2 \xi^{-2}}}{\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q}) - \sqrt{[\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})]^2 + v_{\vec{k}+\vec{Q}}^2 \xi^{-2}}} \right| \quad (7)$$

To see how the shadow band solution appears, let us look at $\Sigma'(\omega, \vec{k})$ at frequencies $|\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})| \gg v_{\vec{k}+\vec{Q}}/\xi$. In this case Eq. (7) reduces to:

$$\Sigma'(\omega, \vec{k}) \approx \frac{g'}{2\pi} \frac{T \ln \xi}{\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})} \quad (8)$$

When $T \ln \xi = \text{const}$ as $T \rightarrow 0$, the large value of $|\Sigma'|$ for $|\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})| \sim v_{\vec{k}+\vec{Q}}/\xi$ leads to the appearance of two solutions for the quasiparticle equation. The effect exists in $D = 2$ in the R. C. regime $\xi \propto \exp(\text{const}/T)$. Since in the weak-to-intermediate coupling regime T_X is of the order of the gap Δ in the ordered state, the condition $\xi \gg \xi_{th} \equiv v_F/T$ can be also written as $\xi \gg \xi_{coh}$, where $\xi_{coh} = \Delta/v_F$ is the coherence length in the ordered state. In three dimensions precursors of AFM are very weak⁶ because the integration of Eq.(3) gives in 3D $\Sigma''(\omega, \vec{k}) \propto g'T \ln \left[[\omega - \tilde{\varepsilon}(\vec{k} + \vec{Q})]^2 + v_{\vec{k}+\vec{Q}}^2 \xi^{-2} \right]$, which is much less singular than in two-dimensional case Eq.(6). In the dimensions $D > 3$ the precursors of AFM bands do not appear even at the very point of the phase transition, because of the large phase space in Eqs.(1),(3). This result could be expected on general physical grounds, since in three and larger dimensions the gap is equal to zero at the Néel temperature. In contrast, in two-dimensions classical thermal fluctuations suppress long-range order at finite temperature and at the $T = 0$ phase transition the system goes directly into the ordered state with a *finite* gap. Finally, I note that the one-dimensional case is, as usual, very special. For the most recent studies of shadow bands in the 1D case see^{14,15}.

IV. CONCLUSIONS

I have shown that in 2D there are a few different antiferromagnetic precursor effects in the spectral function $A(\omega, \vec{k})$, which are realized under different physical conditions. The conditions for dispersive shadow features and shadow bands are given by $\omega_{sf} \ll T$ and $\xi \gg \xi_{th} \equiv v_F/T$. The first conditions insures that the

short-range AFM order is *quasi-static*, which means that not only the susceptibility $\chi'_{sp}(0, \vec{q})$ has a Lorentzian form at $q \approx Q$, but also the static structure factor $S_{sp}(\vec{q})$ has this form. The second condition $\xi \gg \xi_{th} \equiv v_F/T$ implies that the AFM correlation length should be compared with the thermal wave length $\xi_{th} = v_F/T$ of electrons, rather than with the lattice constant a , as was assumed in some previous works on shadow bands. The above conditions for dispersive shadow bands can be realized in two-dimensions when at $T = 0$ there is long-range AFM order. The situation is favorable in two dimension because this is the lower critical dimension for the AFM instability and the mean field phase transition is replaced by the crossover T_X to the regime with very rapidly growing correlation length $\xi \propto \exp(const/T) \gg v_F/T$ for $T \rightarrow 0$.

The dispersive shadow features and shadow bands are closely related to one another. When the correlation length starts to increase faster than $\xi_{th} = v/T$ ($1/T^{1/2}$ at Van Hove point) the shadow bands appear first at the hot spots, while shadow features appear at the wave vectors $\vec{k}$ away from the Fermi surface. When the correlation length increases further the shadow feature is eventually replaced by the shadow band. In this context, the thermally excited shadow feature could be considered as a precursor of the shadow band.

The theory predicts no dispersive shadow features or shadow bands at zero temperature no matter how large is the correlation length. I note that the reason for the absence of dispersive shadow peaks at $T = 0$ is similar to the reason for the absence of shadow peaks in higher dimensions, namely, large phase space. Indeed, in the Matsubara formalism the imaginary axis plays the role of an additional dimension at $T = 0$, since $T \sum_{\omega_n} (...) \rightarrow \int (...) d(i\omega)/2\pi$.

In addition to the dispersive shadow peaks which exist in both $\vec{k}$ and ω scans of the spectral function, there is also a non-dispersive shadow feature¹⁰ in the k -scan only of $A(\omega_{Fixed}, \vec{k})$. The latter effect is due to the *dynamical* quantum spin fluctuation and it exists at zero temperature and finite frequency if ξ is sufficiently large.

Experimental implications: Close to optimal doping the increase of ξ with temperature is well described by the mean-field Curie-Weiss law $\xi \propto (A + BT)^{-1/2}$, i. e. slower than $1/T^{1/2}$ ¹¹. Under those conditions, the present theory does not predict any dispersive shadow feature or shadow band. I also found that the non-dispersive shadow feature due to quantum fluctuations¹⁰ in $A(\omega_{Fixed}, \vec{k})$ is always a very small effect. I did not find such shadow feature for the conditions of the type II experiment of Aebi et al.¹. I conclude that, based on the present approach, the “shadow bands” observed in B2212 close to optimal doping^{1,2} cannot be explained as precursors of AFM bands. The structural explanation of the “shadow bands” in optimally doped materials, have been suggested in Ref.^{4,16} (see also comment¹⁷).

I suggest that precursors of AFM bands should be

searched for in strongly underdoped materials in the RC regime $\xi \propto \exp(const/T)$. These bands should develop first at the crossing points $\vec{k} = \vec{k}_{cr}$ and only much later along the diagonal direction $(0, 0) - (\pi, \pi)$.

I am grateful to M. R. Norman and A. M.-S. Tremblay for many suggestions and stimulating discussions. I also acknowledge useful discussions with A. V. Chubukov and J. Schmalian. This work was supported by the National Science Foundation (Grant No. NSF-DMR-91-20000) through the Science and Technology Center for Superconductivity.

-
- ¹ P. Aebi et al., Phys. Rev. Lett. **72**, 2757 (1994).
 - ² S La Rosa et al., preprint cond-mat/9512132.
 - ³ A.P. Kampf, and J.R. Schrieffer, Phys. Rev. B **41**, 6399 (1990); Phys. Rev. B **42**, 7967 (1990).
 - ⁴ S. Chakravarty, Phys. Rev. Lett. **74**, 1885 (1995).
 - ⁵ P. Aebi et al., J. of Elect. Spectroscopy, **76**, 127 (1995).
 - ⁶ Y. M. Vilks and A.-M. S. Tremblay, Europhys. Lett., **33**, p 159 (1996); preprint cond-mat/9502015.
 - ⁷ J. J. Deisz, D. W. Hess, J. W. Serene, Phys. Rev. Lett. **76**, 1312 (1996).
 - ⁸ M. Langer, J. Schmalian, S. Grabowski, and K.H. Bennemann, Phys. Rev. Lett. **75**, 4508 (1995)
 - ⁹ S. Haas, A. Moreo, and E. Dagotto, Phys. Rev. Lett. **74**, 4281 (1995).
 - ¹⁰ A. V. Chubukov, Phys. Rev. B, **52**, R3847 (1995).
 - ¹¹ A. J. Millis, H. Monien, and D. Pines, Phys. Rev. B. **42**, 167 (1990); P. Monthoux and D. Pines, ibid **47**, 6069 (1993).
 - ¹² The value of the chemical potential for the interacting system in this case happens to be almost the same as the non-interacting one $\mu_0 = -1.31t$.
 - ¹³ Similar expression for Σ have been used by P. A. Lee, T. M. Rice, and P. W. Anderson, Phys. Rev. Lett. **31**, 462 (1973) to study the effect of classical charge fluctuations in quasi-1D electron system.
 - ¹⁴ R. H. McKenzie and D. Scaratt, preprint cond-mat/9609179; Phys. Rev. B, November, 1996.
 - ¹⁵ K. Penc, K. Hallberg, F. Mila and H. Shiba, preprint cond-mat/9512152.
 - ¹⁶ D. J. Singh and W. E. Pickett, Phys. Rev. B **51**, 3128 (1995).
 - ¹⁷ The authors of Ref.² argued against the structural explanation of their experiment because they found only small ω dependence of the amplitude of the shadow peak. However, the amplitude of the weak shadow peak is very sensitive to the used “background” subtraction procedure. Clear description and justification of such procedure, which is absent in Ref.², is necessary to obtain a definite conclusion on this issue.

FIG. 1. The ω -dependence of $A(\omega, \vec{k})$ for $\vec{k} = k_{cr} = (0.825, 0.175)\pi$. The inset show the F. S. (filled line) and the shadow F. S. (dashed line).

FIG. 2. The ω -dependence of $f(\omega)A(\omega, \vec{k})$ (upper panel) and $\Sigma''(\omega, \vec{k})$ (lower panel) for $\vec{k} = (0.66, 0.66)\pi$, that is slightly inside Sh. F. S. The inset shows the $\vec{k}$ -dependence of $A(\omega, \vec{k})$ in the vicinity of $k_{Sh.F.S} = (0.63, 0.63)\pi$ for $\omega = -0.03t$ (∇) and $\omega = -0.125t$ ($\square$).



